

RHEUMATOID ARTHRITIS

Response to comment on “*Aggregatibacter actinomycetemcomitans*–induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis”

Maximilian F. König,¹ Jon T. Giles,² Ricardo P. Teles,^{3,4*} Niki M. Moutsopoulos,⁵ Felipe Andrade^{6†}

Antibodies to leukotoxin A are markers that link *Aggregatibacter actinomycetemcomitans*–associated periodontitis and rheumatoid arthritis.

We thank Volkov and colleagues for their interest in our study and attempt to replicate part of our findings. In our study, we provided mechanistic and clinical evidence that implicates the periodontal pathogen *Aggregatibacter actinomycetemcomitans* (hereinafter *Aa*) as a candidate trigger of autoimmunity in rheumatoid arthritis (RA) (1). We showed that among different periodontal pathogens, *Aa* has the ability to dysregulate peptidylarginine deiminases in neutrophils and induce patterns of hypercitrullination found in periodontitis and the rheumatoid joint. We demonstrated that *Aa*-induced hypercitrullination is mediated by its major virulence factor leukotoxin A (LtxA). Using antibodies to LtxA as surrogate markers of *Aa* infection, we found that exposure to leukotoxic *Aa* strains was highly prevalent both in periodontitis and in RA when compared with individuals without periodontitis. Exposure to *Aa* was associated with distinct anti-citrullinated protein antibody (ACPA) specificities in RA, strengthening the hypothesis that *Aa*-induced hypercitrullination may be relevant to RA pathogenesis by inducing ACPAs. Moreover, a significant association of human leukocyte antigen–DRB1 shared epitope (SE) alleles with ACPA and rheumatoid factor (RF) positivity was only observed in RA patients exposed to *Aa* [P values for interaction: $P = 0.022$ for ACPA positivity, $P = 0.022$ for anti-cyclic citrullinated peptide 2 (CCP2), and $P = 0.012$ for RF] (1). Together, these studies provide a mechanism to explain the tantalizing hypothesis that periodontitis is a driver of RA (1).

In their comment, Volkov *et al.* (2) only addressed the clinical associations reported in our study. First, they asked whether exposure to *Aa* as measured by anti-LtxA antibodies is specific for RA. Second, the authors attempted to replicate associations between *Aa* exposure, SE alleles, and ACPAs in a group of 594 arthritis patients selected from the Leiden Early Arthritis Clinic (EAC) cohort. Their analysis highlights important differences in data interpretation and cohorts studied. Despite this, we believe that their data are consistent with our findings.

¹Department of Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA. ²Division of Rheumatology, Columbia University, College of Physicians and Surgeons, New York, NY 10032, USA. ³Department of Periodontology, School of Dentistry, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA. ⁴Department of Applied Oral Sciences, Forsyth Institute, Cambridge, MA 02142, USA. ⁵National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD 20892, USA. ⁶Division of Rheumatology, The Johns Hopkins University School of Medicine, Baltimore, MD 21224, USA.

*Deceased.

†Corresponding author. Email: andrade@jhmi.edu

Different from the authors' assumption, our study did not suggest specificity of *Aa* exposure to RA. *Aa* has been firmly established as a pathogen associated with periodontitis (3–5). A considerable portion of our manuscript is dedicated to characterizing anti-LtxA antibodies as markers of periodontal disease status [figure 6B; figure S4, A and B; and table S4 of (1)], and we showed that anti-LtxA concentrations correlate with periodontitis severity in patients without RA [$\beta = 1.12$; $P < 0.0001$; figure S4B of (1)]. The idea that antibodies to LtxA may be restricted to patients with RA is therefore rejected by our data and misinterpreted by Volkov *et al.* (2). Periodontitis is a chronic inflammatory disease that is highly prevalent in the general population. The presence of antibodies against LtxA in any individual with periodontitis exposed to *Aa* (with or without any chronic illness) is therefore expected. Regardless of these data, Volkov *et al.* initially demonstrated that antibodies to LtxA are associated with *Aa* infection in periodontitis patients without RA [figure 1A of (2)], which confirmed our data. Having again demonstrated that anti-LtxA antibodies are markers of *Aa*-associated periodontitis in individuals without RA, the authors tested whether these antibodies are exclusively found in RA compared to healthy individuals or patients with other arthritides of unknown periodontal status.

Different from our study, Volkov *et al.* (2) did not include healthy controls without periodontitis to define cutoffs for anti-LtxA positivity. In the absence of clinically relevant cutoffs to define exposure to leukotoxic strains of *Aa*, it is impossible to identify individuals who are truly positive for anti-LtxA antibodies. Using an arbitrary cutoff instead (“the lowest point of the linear part of the standard curve” of mixed RA patient sera), Volkov *et al.* (2) reported that 58% of patients with RA have antibodies against LtxA. Although this exceeds the prevalence of anti-LtxA positivity in our cohort (43%) (1), it supports the high prevalence of *Aa* exposure in patients with RA. The authors further showed that anti-LtxA antibodies were significantly enriched in RA as compared to healthy controls ($P < 0.0001$) and patients with other inflammatory arthritides ($P < 0.001$ for psoriatic arthritis and $P < 0.05$ for spondyloarthritis with peripheral arthritis). Moreover, anti-LtxA antibody concentrations in RA were markedly higher than in any other form of arthritis, although sample size appeared prohibitive to reach statistical significance in some groups [figure 1B of (2)]. The presence of anti-LtxA positivity reported by Volkov *et al.* in healthy controls and patients with non-RA arthritides likely reflects that some of these individuals have *Aa* infection and periodontitis, which is an oral pathology that is not exclusively found in RA. Regarding the consequences of *Aa* exposure, however, our model

Copyright © 2018
The Authors, some
rights reserved;
exclusive licensee
American Association
for the Advancement
of Science. No claim
to original U.S.
Government Works

proposes that only individuals at risk for RA (through the carriage of genetic susceptibility genes) will develop autoimmune sequelae from *Aa*-induced hypercitrullination (1). It is therefore expected that not everybody who is infected with *Aa* will develop RA.

Although absolute numbers of positivity for anti-LtxA reported by the authors in figure 1B of (2) are impossible to ascertain in the absence of non-periodontitis controls, a more stringent cutoff for positivity may have been more consistent with our own validated data set. Regardless of these limitations, the association of anti-LtxA antibodies with RA in the EAC cohort is compelling. Yet, the implications of these results in the context of our work were unfortunately not discussed by the authors.

Altogether, we believe that these data strongly support our findings by (i) validating anti-LtxA antibodies as powerful tools to identify individuals exposed to *Aa*, (ii) reemphasizing the association of anti-LtxA antibodies with chronic periodontitis, and (iii) confirming a high prevalence and significant enrichment of *Aa* exposure in early RA as compared to controls and other inflammatory arthritides. The authors' assertion that patients with RA may be "more frequently anti-LtxA-positive simply because they have more periodontitis" is epidemiologically correct, but in no way negates an etiologic role of *Aa* in the genetically susceptible host.

Volkov *et al.* (2) next analyzed the association of anti-LtxA antibodies with anti-CCP positivity (referred to as "ACPA" in their comment) and SE alleles in RA. The authors showed that anti-LtxA antibody titers do not differ by either anti-CCP or SE status in RA. Similar to their findings, we reported that anti-LtxA positivity was not associated with anti-CCP antibodies [table S5 of (1)]. In this respect, their findings are consistent with our own data. In our study, anti-LtxA positivity was only significantly associated with distinct ACPA fine specificities measured by Bio-Plex bead-based assays [citrullinated antigens: apolipoprotein A1, $P < 0.001$; apolipoprotein E, $P = 0.004$; histone H2B, $P = 0.020$; vimentin, $P = 0.018$; apolipoprotein E (277–296, cyclic peptide), $P = 0.032$; clusterin (231–250, cyclic peptide), $P = 0.020$; and hnRNP B1b (RA33), $P = 0.038$; table 1 of (1)], but not with anti-CCP positivity in general [table S5 of (1)]. We showed that the known association of SE alleles and autoantibody positivity (anti-CCP, Bio-Plex bead-based ACPA, or RF) was restricted to RA patients with *Aa* exposure [table 2 of (1)]. In their analysis (which only included anti-CCP), Volkov *et al.* found only a small increase in the interaction of anti-LtxA and SE alleles for anti-CCP positivity in the Leiden cohort.

Several important differences in cohorts and methodology may account for this difference in effect size: (i) The authors' cohort is known to differ substantially from our cohort in anti-CCP positivity (only 58% in Leiden versus 77% in Baltimore) (1). These differences are not easily explained by ethnicity because the authors reported ACPA positivity in 75% of Dutch patients with established RA as part of the IMPROVED (Induction therapy with Methotrexate and Prednisone in Rheumatoid Or Very Early arthritic Disease) cohort (these patients fulfilled the American College of Rheumatology 1987 criteria for RA at the time of inclusion) (6). Compared with other cohorts, there is a marked discrepancy between the expected and reported anti-CCP positivity in the EAC cohort. This was acknowledged in a recent study in which the authors reported the frequency of anti-CCP2 positive RA as only 51% in the EAC (total cohort) as compared with 65% in IMPROVED. Both frequencies are lower than that in our cohort [77% in ESCAPE RA (Evaluation of Subclinical Cardiovascular disease And Predictors of Events in RA)], and this was suggested to be due to differences in inclusion criteria by the authors. Notably, the Leiden

EAC includes patients with recent onset of arthritis in whom definitive diagnoses are established after 1 year of follow-up (6). The large subset of anti-CCP-negative RA in the EAC cohort may therefore be composed of early arthritis patients who were either misclassified as having RA (7, 8), or who will have milder forms of disease. Notably, anti-peptidylcitrulline reactivities in the Leiden EAC were observed in a larger subset of RA patients by multiplex array (64%) when compared with anti-CCP2 (51%) (6), suggesting that in this early cohort, CCP2 may not be sensitive enough to capture the entirety of the evolving ACPA repertoire. Moreover, given the range of disease durations in our cohort (from early disease to many decades), which is wider than that of the Leiden EAC cohort, a potential scenario for interpreting our finding is that the context of *Aa* may be required to observe the evolution of the ACPA repertoire, as well as other autoantibodies, among those with SE over the longer course of clinical disease, and not just in the earliest phase. Certainly, further studies are required to confirm this hypothesis. (ii) The authors' definitions of anti-LtxA positivity differ considerably from our cohort (cutoffs were selected arbitrarily by Volkov *et al.*, but experimentally established using nonperiodontitis healthy controls in our study). Despite not affecting comparisons of median antibody concentrations between groups, uncertainty in cutoff selection can alter any interaction analyses of anti-CCP, SE, and anti-LtxA status. (iii) Differences in racial and ethnical composition of our cohort (Baltimore) as compared with Europe (Leiden) may have influenced the effect of *Aa* in inducing ACPAs, although this is less likely.

In summary, our study provides mechanistic insights into how the periodontal pathogen *Aa* promotes hypercitrullination and neutrophil cell death, suggesting a biologically plausible link between chronic periodontal infection and the promotion of autoimmunity directed against citrullinated proteins. Virulent strains of *Aa* are strongly associated with chronic and aggressive periodontitis, and may explain the intriguing association of RA and periodontitis in a subgroup of patients with RA. Notably, *Aa* is not a pathogen exclusive to RA. The identification of a single bacterial species in RA that can fulfill Henle (9) and Koch's (10) postulates of causation, as implied by Volkov *et al.* (2), is highly unlikely. Instead, *Aa*-induced hypercitrullination provides a model to understand how pathobionts may influence the development of autoimmune diseases in susceptible individuals. It is certain that further studies are necessary to confirm our model. However, we believe that the interpretation of the data and conclusions presented by Volkov *et al.* cannot exclude a biologically relevant role for *Aa* in the etiopathogenesis of RA.

REFERENCES AND NOTES

1. M. F. König, L. Abusleme, J. Reinholdt, R. J. Palmer, R. P. Teles, K. Sampson, A. Rosen, P. A. Nigrovic, J. Sokolove, J. T. Giles, N. M. Moutsopoulos, F. Andrade, *Aggregatibacter actinomycetemcomitans*-induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis. *Sci. Transl. Med.* **8**, 369ra176 (2016).
2. M. Volkov, J. Dekkers, B. G. Loos, S. Bizzarro, T. W. J. Huizinga, H. A. Praetorius, R. E. M. Toes, D. van der Woude, Comment on "Aggregatibacter actinomycetemcomitans-induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis". *Sci. Transl. Med.* **10**, eaan8349 (2018).
3. C. Chen, T. Wang, W. Chen, Occurrence of *Aggregatibacter actinomycetemcomitans* serotypes in subgingival plaque from United States subjects. *Mol. Oral Microbiol.* **25**, 207–214 (2010).
4. N. Pahumunto, P. Ruangsri, M. Wongsuwanlert, S. Piwat, G. Dahlen, R. Teanpaisan, *Aggregatibacter actinomycetemcomitans* serotypes and DGGE subtypes in Thai adults with chronic periodontitis. *Arch. Oral Biol.* **60**, 1789–1796 (2015).
5. M. Minguez, X. Pousa, D. Herrera, A. Blasi, M. C. Sánchez, R. León, M. Sanz, Characterization and serotype distribution of *Aggregatibacter actinomycetemcomitans* isolated from a population of periodontitis patients in Spain. *Arch. Oral Biol.* **59**, 1359–1367 (2014).

6. J. van Heemst, L. A. Trouw, L. Nogueira, H. W. van Steenberghe, A. H. M. van der Helm-van Mil, C. F. Allaart, G. Serre, R. Holmdahl, T. W. J. Huizinga, R. E. M. Toes, D. van der Woude, An investigation of the added value of an ACPA multiplex assay in an early rheumatoid arthritis setting. *Arthritis Res. Ther.* **17**, 276 (2015).
7. Y. Kaneko, M. Kuwana, H. Kameda, T. Takeuchi, Sensitivity and specificity of 2010 rheumatoid arthritis classification criteria. *Rheumatology (Oxford)* **50**, 1268–1274 (2011).
8. H. Radner, T. Neogi, J. S. Smolen, D. Aletaha, Performance of the 2010 ACR/EULAR classification criteria for rheumatoid arthritis: A systematic literature review. *Ann. Rheum. Dis.* **73**, 114–123 (2014).
9. J. Henle, *Von den Miasmen und Kontagien: Und von den miasmatisch-kontagiösen Krankheiten (1840)* (J.A. Barth, 1910).
10. R. Koch, Über bakteriologische Forschung, in *Verhandlungen des X. Internationalen Medicinischen Congresses, Berlin* (August Hirschwald Verlag, 1890), vol. 1.

Funding: F.A. was supported by the Jerome L. Greene Foundation and National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS)/NIH grant R01AR069569. R.P.T. was supported by National Institute of Dental and Craniofacial Research (NIDCR)/NIH grant

DE021127-01. N.M.M. was supported by the Intramural Research Program of the NIDCR/NIH. ESCAPE RA was supported by NIAMS/NIH AR050026-01. The content is solely the responsibility of the authors and does not necessarily represent the official views of NIAMS or the NIH.

Competing interests: F.A. is an inventor on issued patent no. 8,975,033 held by The Johns Hopkins University that covers "Human autoantibodies specific for PAD3 which are cross-reactive with PAD4 and their use in the diagnosis and treatment of rheumatoid arthritis and related diseases." F.A. received a grant from MedImmune and has served as consultant for Bristol-Myers Squibb Company and Pfizer.

Submitted 5 July 2017

Accepted 20 February 2018

Published 21 March 2018

10.1126/scitranslmed.aao3031

Citation: M. F. Konig, J. T. Giles, R. P. Teles, N. M. Moutsopoulos, F. Andrade, Response to comment on "Aggregatibacter actinomycetemcomitans-induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis." *Sci. Transl. Med.* **10**, eaao3031 (2018).

Science Translational Medicine

Response to comment on "*Aggregatibacter actinomycetemcomitans*–induced hypercitrullination links periodontal infection to autoimmunity in rheumatoid arthritis"

Maximilian F. König, Jon T. Giles, Ricardo P. Teles, Niki M. Moutsopoulos and Felipe Andrade

Sci Transl Med **10**, eaao3031.
DOI: 10.1126/scitranslmed.aao3031

ARTICLE TOOLS	http://stm.sciencemag.org/content/10/433/eaao3031
RELATED CONTENT	http://stm.sciencemag.org/content/scitransmed/10/433/eaan8349.full http://stm.sciencemag.org/content/scitransmed/8/369/369ra176.full http://stm.sciencemag.org/content/scitransmed/8/331/331ra38.full http://stm.sciencemag.org/content/scitransmed/7/290/290ra87.full
REFERENCES	This article cites 8 articles, 2 of which you can access for free http://stm.sciencemag.org/content/10/433/eaao3031#BIBL
PERMISSIONS	http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the [Terms of Service](#)

Science Translational Medicine (ISSN 1946-6242) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science Translational Medicine* is a registered trademark of AAAS.