

Supplementary Materials for

Phytosterols Promote Liver Injury and Kupffer Cell Activation in Parenteral Nutrition–Associated Liver Disease

Karim C. El Kasmi, Aimee L. Anderson, Michael W. Devereaux, Padade M. Vue, Wujuan Zhang, Kenneth D. R. Setchell, Saul J. Karpen, Ronald J. Sokol*

*Corresponding author. E-mail: Ronald.Sokol@childrenscolorado.org

Published 9 October 2013, *Sci. Transl. Med.* **5**, 206ra137 (2013)

DOI: 10.1126/scitranslmed.3006898

This PDF file includes:

Fig. S1. Experimental design and experimental treatment groups.

Other Supplementary Material for this manuscript includes the following:

(available at

www.sciencetranslationalmedicine.org/cgi/content/full/5/206/206ra137/DC1)

Table S1 (Microsoft Excel format). Original data for all figures.

Table S2 (Microsoft Excel format). Statistical analyses for all figures

Fig. S1. Experimental design and experimental treatment groups.




Fig. S1 Legend. Experimental design and experimental treatment groups.

Blood and tissue were collected from all mice at the end of the experiment for measurement of serum AST, ALT, bilirubin, bile acids; liver macrophage isolation; liver and macrophage qPCR.